

REGULARITY AND EXTREMALITY OF QUASICONFORMAL HOMEOMORPHISMS ON CR 3-MANIFOLDS

PUQI TANG

ABSTRACT. This paper first studies the regularity of conformal homeomorphisms on smooth locally embeddable strongly pseudoconvex CR manifolds. Then moduli of curve families are used to estimate the maximal dilatations of quasiconformal homeomorphisms. On certain CR 3-manifolds, namely, CR circle bundles over flat tori, extremal quasiconformal homeomorphisms in some homotopy classes are constructed. These extremal mappings have similar behaviors to Teichmüller mappings on Riemann surfaces.

1. INTRODUCTION

A contact manifold M is a manifold of odd dimension with a non-integrable distribution HM of tangent hyperplanes. A Cauchy-Riemann (CR) manifold is a contact manifold M endowed a complex structure on the contact bundle HM . Two CR manifolds are equivalent if there is a homeomorphism between them which preserves both contact and CR structures. Generally, between any two CR structures assigned on the same contact manifold, there may be no such so-called CR homeomorphism between them. Therefore we consider those homeomorphisms between CR manifolds which preserve the underlying contact structures and distort the CR structures boundedly. They are called quasiconformal homeomorphisms. In a class of homeomorphisms between two CR manifolds, an extremal mapping is a quasiconformal homeomorphisms which distorts the CR structures in a minimal way. This paper studies regularity of quasiconformal homeomorphisms and extremal quasiconformal homeomorphisms on smooth strongly pseudoconvex CR manifolds.

The notion of quasiconformal homeomorphisms is a new tool to study CR structures as initiated by Korányi and Reimann. In this paper, we use an analytic definition of quasiconformal homeomorphisms given in [12] which is a generalization of the one given by Korányi and Reimann in [5]. We restrict ourselves to the 3-dimensional case here not only because the notion of quasiconformality is not invariant under CR transformations in higher dimensional cases, but also because 3-dimensional CR structures are among the most interesting objects in the theory of CR manifolds. We refer to [7] for details about the second point.

1991 *Mathematics Subject Classification.* 32G07.

Key words and phrases. CR manifold, quasiconformal homeomorphism, modulus of a curve family, Legendrian foliation, sub-Riemannian geometry, Teichmüller mapping.

Korányi and Reimann proved that C^4 conformal homeomorphisms on Heisenberg groups must be smooth and CR (Theorem 8, [6]). By applying a regularity theorem of weak CR mappings of Pinchuk and Tsyganov, we generalize Korányi and Reimann’s result to that a conformal homeomorphism f between two smooth, strongly pseudoconvex, locally embeddable CR manifolds must be smooth and CR, if f has L^1_{loc} horizontal derivatives (Theorem 2.3). Hence between such CR manifolds, quasiconformal homeomorphisms with this weak regularity are actually “quasi-CR”.

To study the extremality of quasiconformal homeomorphisms is a global problem. But our analytic definition of quasiconformality is proposed infinitesimally. Therefore we need some global notion to describe the quasiconformality. The one best fitting our later developments is the notion of moduli of curve families. We prove that a C^2 diffeomorphism is quasiconformal if and only if it preserves moduli of certain curve families up to a fixed bounded multiple (Theorem 3.3). On the other hand, a homeomorphism satisfying this property is absolutely continuous on lines (ACL) (Theorem 3.4).

Between CR circle bundles over flat tori, we construct extremal quasiconformal homeomorphisms in certain homotopy classes (Theorem 4.2). There are two transversal Legendrian foliations such that the extremal homeomorphism constructed preserves these two foliations. More precisely, it is a stretching by a constant factor along leaves of one foliation and a compressing by the same factor along leaves of another foliation. This behavior is analogous to those of Teichmüller mappings on Riemann surfaces (see [13] or [1]). The generator T of the circle action is transversal to the contact bundle. In this transversal direction, the extremal mappings are equivariant under the circle action.

But on an arbitrary CR 3-manifold with a transversal free circle action, an extremal quasiconformal homeomorphism is not necessarily equivariant under the circle action. Such CR manifolds are constructed in [12] so that no extremal quasiconformal mapping between them is equivariant.

This work is heavily influenced by the theory of quasiconformal homeomorphisms on Riemann surfaces and Teichmüller theory. Numerous proofs in this paper are motivated by the proofs of the analogous facts on Riemann surfaces. For example, the construction of the extremal homeomorphisms made in Theorem 4.2, one of the main results of this paper, can find its root in the classical Grötzsch’s theorem which is proved by a length-area argument [9]. Teichmüller generalized this result to closed Riemann surfaces, in particular tori, by an ergodic version of the length-area argument [13]. The notion of modulus of a curve family is a formalism of the length-area (volume) argument. In section 4, we reformulate Teichmüller’s method to the CR setting by computing moduli of some special families of curves and successfully find the extremal quasiconformal homeomorphisms in certain homotopy classes of mappings between CR circle bundles over flat tori.

Acknowledgement. This author is very grateful to his academic advisor, László Lempert for the guidance with great insights. Thanks also to David Drasin and Juha Heinonen for helpful talks.

2. REGULARITY OF CONFORMAL HOMEOMORPHISMS

For $j = 1, 2$, let M_j be a smooth strongly pseudoconvex CR 3-manifolds. The contact

bundles HM_j is assumed to be smooth and orientable, that is, there exists smooth global 1-form η_j on M_j , which is called a contact form, so that $HM_j = \text{Ker } \eta_j$. Let $J_j : HM_j \rightarrow HM_j$ denote the CR structure on M_j . $H^{0,1}M_j \triangleq \{X + iJ_jX \mid X \in HM_j\} \subset \mathbb{C} \otimes HM_j$ is the $(0,1)$ tangent bundle on M_j . $\wedge^{1,0}M_j \triangleq \{\text{linear functional } \psi : \mathbb{C} \otimes HM_j \rightarrow \mathbb{C} \mid \psi(J_jX) = i\psi(X), \text{ for } X \in HM_j\}$.

A mapping $f : M_1 \rightarrow M_2$ is said to be absolutely continuous on lines (ACL) if for any open set with a smooth contact fibration, f is absolutely continuous along all fibers in this fibration except a subfamily of measure zero. Here a subfamily of fibers of the fibration is said to have measure zero if intersections of these fibers with any transversal regular surface has measure zero on the surface (see [12]).

Definition 2.1. A homeomorphism $f : M_1 \rightarrow M_2$ is said to be K -quasiconformal for a finite constant $K \geq 1$ if

- (i) f is ACL;
- (ii) f is differentiable almost everywhere, and its differential f_* preserves the contact structures, i.e., $f_*(H_qM_1) \subset H_{f(q)}M_2$, for $q \in M_1$ where f has differential.
- (iii) for norms $|\cdot|_1$ and $|\cdot|_2$ defined by any Hermitian metrics on HM_1 and HM_2 respectively,

$$(2.1) \quad K(f)(q) = \frac{\max_{X \in H_qM_1, |X|_1=1} |f_*(X)|_2}{\min_{X \in H_qM_1, |X|_1=1} |f_*(X)|_2} \leq K < \infty,$$

for almost all $q \in M_1$. Here $K = \text{ess sup}_{q \in M_1} K(f)(q)$ is called the maximal dilatation of f . f is conformal if $K(f) = 1$. $K(f) = \infty$ if f is not K -quasiconformal for any finite $K \geq 1$.

A mapping $f : M_1 \rightarrow M_2$ which is differentiable almost everywhere is said to have L_{loc}^p horizontal derivatives for $p \geq 1$ if for any smooth function $h : M_2 \rightarrow \mathbb{R}$ and any smooth local section X of HM_1 on an open set $U \subset\subset M_1$, the function $X(h \circ f)$ which is defined almost everywhere on U is in $L^p(U)$.

A mapping $f : M_1 \rightarrow M_2$ is said to have L_{loc}^p weak horizontal derivatives for some $p \geq 1$ if for any smooth function $h : M_2 \rightarrow \mathbb{R}$ and any open set $U \subset\subset M_1$ with a smooth local section $X \neq 0$ of HM_1 on U , there exists a function $g \in L_{\text{loc}}^p(M_1)$ so that

$$(2.2) \quad \int_U X\phi \cdot (h \circ f) dv_1 = - \int_U g\phi dv_1$$

for all $\phi \in C_0^\infty(U)$, where dv_1 is a smooth volume form on M_1 . Certainly, the function g depends on the choice of dv_1 .

For the proof of the next theorem, we fix a norm $|\cdot|$ on HM_1 . A regular curve on a contact manifold is called Legendrian if it is tangent to the contact structure. For any smooth Legendrian curve $\gamma : I \rightarrow M_1$ with an interval $I \in \mathbb{R}$ and a function g defined on an open neighborhood of γ , define the line integral

$$\int_\gamma g = \int_I g(\gamma(t)) |\gamma'(t)| dt.$$

Theorem 2.2. *A mapping $f : M_1 \rightarrow M_2$ is ACL and has L^p_{loc} horizontal derivatives for some $p \geq 1$ if and only if f has L^p_{loc} weak horizontal derivatives.*

Proof. First assume that the homeomorphism $f : M_1 \rightarrow M_2$ is ACL and has L^p_{loc} horizontal derivatives. Let $h : M_2 \rightarrow \mathbb{R}$ be a smooth function and $U \subset\subset M_1$ be any open set with a smooth section $X \neq 0$ of HM_1 on it.

We can assume that the trajectories $\Gamma = \{\gamma\}$ of X form a contact fibration of U by shrinking U appropriately. Let Γ_1 be the subfamily of those $\gamma \in \Gamma$ along which f is absolutely continuous. Then $\Gamma \setminus \Gamma_1$ has measure zero. Along $\gamma \in \Gamma_1$, $h \circ f$ is absolutely continuous, so $X(h \circ f)$ exists almost everywhere on γ and

$$(2.3) \quad \int_{\gamma} X\phi \cdot (h \circ f) + \int_{\gamma} \phi \cdot X(h \circ f) = \int_{\gamma} X(\phi \cdot h \circ f) = 0, \quad \forall \phi \in C_0^\infty(U).$$

We have topological and differential structures on Γ such that the natural projection $p : U \rightarrow \Gamma$ is open and smooth. Then Γ becomes a smooth surface. Let t be the parameter of the flow generated by X and ω be any area form on Γ . Then $dv_1 \triangleq p^*\omega \wedge dt$ is a volume form of U . Integrating the expressions in (2.3) against ω with respect to $\gamma \in \Gamma_1$, then by the ACL property of f , local L^p integrability of $X(h \circ f)$ and Fubini's theorem, we obtain

$$(2.4) \quad \int_U X\phi \cdot (h \circ f) dv_1 = - \int_U \phi \cdot X(h \circ f) dv_1.$$

So the weak derivative of $h \circ f$ in the X direction is given by $X(h \circ f) \in L^p_{\text{loc}}$.

Conversely, assume f has L^p_{loc} weak horizontal derivatives. For any open set $U \subset\subset M_1$ and a smooth contact fibration Γ of U , let X be the nonzero horizontal vector field on U so that X_q , for any $q \in U$, is the tangent vector at q of the fiber $\gamma \in \Gamma$ passing through q . Let $B \subset\subset U$ be an open set with the coordinate system $\{(x, y, t) \mid a_1 < x < a_2, b_1 < y < b_2, c_1 < t < c_2\}$, here t is the parameter of the flow generated by X , i.e., $X = \frac{\partial}{\partial t}$.

For any smooth function h on M_1 , $h \circ f$ has $L^p(U)$ weak derivative in the direction X . Denote it by $\psi \in L^p(U)$. Hence there exists a sequence of C^1 functions g_n on B such that g_n converges to $h \circ f$ uniformly in B and Xg_n converges to ψ in $L^p(B)$. Let $B_{x,y,t} = (a_1, x) \times (b_1, y) \times (c_1, t)$, $R_{x,y} = (a_1, x) \times (b_1, y)$.

$$(2.5) \quad \int_{B_{x,y,t}} Xg_n(u, v, w) dudvdw = \int_{R_{x,y}} (g_n(u, v, t) - g_n(u, v, c_1)) dudv.$$

Hence by taking limits, we have

$$(2.6) \quad \int_{B_{x,y,t}} \psi(u, v, w) dudvdw = \int_{R_{x,y}} ((h \circ f)(u, v, t) - (h \circ f)(u, v, c_1)) dudv.$$

Let $\{t_n\}$ be a countable dense set of (c_1, c_2) . (2.6) implies for each t_n , there exists a set $E_n \subset R_{a_2, b_2}$ so that $R_{a_2, b_2} \setminus E_n$ is of measure zero and

$$(2.7) \quad \int_{c_1}^{t_n} \psi(x, y, w) dw = (h \circ f)(x, y, t_n) - (h \circ f)(x, y, c_1), \quad \forall (x, y) \in E_n.$$

Then (2.7) is true for all $(x, y) \in E \triangleq \cap E_n$ and all t_n . Hence by continuity of both sides in t ,

$$(2.8) \quad \int_{c_1}^t \psi(x, y, w) dw = (h \circ f)(x, y, t) - (h \circ f)(x, y, c_1), \quad \forall (x, y) \in E, t \in (c_1, c_2).$$

So $(h \circ f)(x, y, t)$ is absolutely continuous in t for $(x, y) \in E$. Note $R_{a_2, b_2} \setminus E$ has measure zero. So f is ACL since $B \subset\subset U$ is an arbitrary rectangular coordinate chart and $U \subset\subset M_1$ is arbitrary in M_1 . Moreover, $X(h \circ f)$ exists almost everywhere and $X(h \circ f) = \psi \in L^p(U)$ on U . So f has L^p_{loc} horizontal derivatives. $\square$

A homeomorphism $f : M_1 \rightarrow M_2$ is said to be weakly CR if for any smooth CR function $h : M_2 \rightarrow \mathbb{C}$, open set $U \subset\subset M_1$, $\phi \in C_0^\infty(U)$ and $\bar{Z} \in H^{0,1}M_1$, $\int_U (h \circ f) \cdot \bar{Z} \phi = 0$.

Theorem 2.3. *Assume M_1 and M_2 are two smooth, strongly pseudoconvex, locally embeddable CR 3-manifolds, $f : M_1 \rightarrow M_2$ is a conformal homeomorphism with L^1_{loc} horizontal derivatives. Then f is smooth and CR.*

Proof. A simple linear algebra argument shows that at a point $q \in M_1$ where f is differentiable

$$(2.9) \quad K(f)(q) = \frac{1 + |\mu(q)|}{1 - |\mu(q)|} \quad \text{with} \quad |\mu(q)| = \left| \frac{\langle f^* \psi_2, \bar{Z} \rangle}{\langle f^* \psi_2, Z \rangle} \right| (q),$$

for any nonzero $\bar{Z} \in H^{0,1}M_1$ and nonzero $\psi_2 \in \wedge^{1,0}M_2$.

Then $K(f) = 1$ implies that if f is differentiable at $q \in M_1$, $\langle f^* \psi_2, \bar{Z} \rangle = 0$ for any $\bar{Z} \in H^{0,1}_q M_1$ and $\psi_2 \in \wedge^{1,0}_{f(q)} M_2$. In other words, f_* preserves the CR structures at the points where it is defined. Thus $\bar{Z}(h \circ f) = 0$ for any CR function h on M_2 and $\bar{Z} \in H^{0,1}M_2$. Theorem 2.2 says that f is weakly CR. M_1 and M_2 are locally embeddable implies they are locally embeddable into $\mathbb{C}^2$ as hypersurfaces. A theorem of Pinchuk and Tsyganov (Theorem 2, [10]) asserts that such f must be smooth, hence CR. $\square$

3. MODULI OF CURVE FAMILIES

Let M be a smooth, compact, contact 3-manifold. We always assume HM is smooth and oriented. A sub-Riemannian metric on M with respect to HM is a smooth positive definite quadratic form on HM . Fix a sub-Riemannian metric on M with respect to HM momentarily, and denote by $|\cdot|$ the corresponding norm on HM . For general theory of sub-Riemannian geometry, we refer to [11] and [2].

The sub-Riemannian metric on M can be extended to a Riemannian metric on M canonically as follows. Let ω be the oriented area form on HM with respect to the sub-Riemannian metric. Then there exists a unique contact form η so that $d\eta|_{HM} = \omega$. Let T be the characteristic vector field of η , namely, T is the unique vector field satisfying that $T \lrcorner \eta = 1$ and $T \lrcorner d\eta = 0$. Declaring T is a unit vector orthogonal to HM , we obtain a Riemannian metric

which is called the canonical extension of the sub-Riemannian metric. The positive volume form of this Riemannian metric is $dv \triangleq d\eta \wedge \eta$.

A curve $\gamma : I_\gamma \rightarrow M$ with an interval $I_\gamma \subset \mathbb{R}$ is called locally rectifiable if γ is absolutely continuous and $\gamma'(t)$ is tangent to HM for almost all $t \in I_\gamma$. γ is called rectifiable if γ is locally rectifiable and the length

$$(3.1) \quad l(\gamma) \triangleq \int_{I_\gamma} |\gamma'(t)| dt < \infty.$$

We set $l(\gamma) = \infty$ if γ is not rectifiable. For a locally rectifiable curve γ and a non-negative Borel-measurable function σ on M , define the line integral

$$(3.2) \quad \int_\gamma \sigma = \int_{I_\gamma} \sigma(\gamma(t)) |\gamma'(t)| dt.$$

Definition 3.1. Let Γ be a family of curves $\gamma : I_\gamma \rightarrow M$. An admissible measure for Γ is a Borel-measurable function $\sigma : M \rightarrow \mathbb{R}$ so that $\sigma \geq 0$ and $\int_\gamma \sigma \geq 1$, for all locally rectifiable $\gamma \in \Gamma$. Denote the set of admissible measures for Γ by $A(\Gamma)$. The modulus of Γ is defined by

$$(3.3) \quad \text{Mod}_M(\Gamma) = \inf_{\sigma \in A(\Gamma)} \int_M \sigma^4 dv.$$

Remark. (1) It is easy to see that if two sub-Riemannian metrics on M with respect to HM define the same conformal structure on HM , then they give the same value to $\text{Mod}_M(\Gamma)$.

(2) If $\Gamma_r \subset \Gamma$ consisting of all locally rectifiable curves of Γ , then $\text{Mod}_M(\Gamma_r) = \text{Mod}_M(\Gamma)$.

The following proposition shows that modulus, regarded as a measure of (more precisely, locally rectifiable) curve families, generalizes the concept of measure zero used in the definition of ACL property. Thereafter if a property holds for all curves in a family Γ except a subfamily with zero modulus, we say this property is true for almost all curves in Γ .

Proposition 3.2. Let U be an open set of M , Γ a contact fibration of U , $\Gamma_1 \subset \Gamma$. Then Γ_1 has measure zero if and only if $\text{Mod}_M(\Gamma_1) = 0$.

Proof. Without loss of generality, we assume U is a domain of the coordinator system $\{(x, y, t) \mid a_1 < x < a_2, b_1 < y < b_2, c_1 < t < c_2\}$ and $X = \frac{\partial}{\partial t}$ is tangent to Γ . Let $E \subset (a_1, a_2) \times (b_1, b_2)$ so that $\Gamma_1 = \{\text{curves } t \mapsto (x, y, t) \mid (x, y) \in E\}$.

If Γ_1 has measure zero, then $\int_E dx dy = 0$. Notice

$$(3.4) \quad \sigma_0 = \begin{cases} \frac{1}{c_2 - c_1}, & \text{when } (x, y) \in E, t \in (c_1, c_2), \\ 0, & \text{otherwise,} \end{cases} \in A(\Gamma_1).$$

Therefore

$$(3.5) \quad \text{Mod}_M(\Gamma_1) \leq c \int_U \sigma_0^4 dx dy dt = 0,$$

where c is a constant upper bound of the Jacobian J on U with $J dx dy dt = dv$.

If $\text{Mod}_M(\Gamma_1) = 0$, then for any $\sigma \in A(\Gamma_1)$ with $\sigma = 0$ outside U , $\gamma \in \Gamma_1$,

$$(3.6) \quad 1 \leq \left(\int_{\gamma} \sigma \right)^4 \leq \left(\int_{\gamma} \sigma^4 \right) \left(\int_{\gamma} 1 \right)^3 = (c_2 - c_1)^3 \int_{\gamma} \sigma^4.$$

Taking the integral over E ,

$$(3.7) \quad \int_E dx dy \leq c' (c_2 - c_1)^3 \int_M \sigma^4.$$

Hence

$$(3.8) \quad \int_E dx dy \leq c' (c_2 - c_1)^3 \text{Mod}_M(\Gamma_1) = 0,$$

that is, Γ_1 has measure zero. $\square$

Let M_1 and M_2 be two compact, smooth, strongly pseudoconvex CR 3-manifolds with smooth contact form η_1 and η_2 respectively. Here the roles of sub-Riemannian metrics on HM_1 and HM_2 are played by Hermitian metrics with respect to the CR structures J_1 and J_2 respectively.

Theorem 3.3. *A C^2 homeomorphism $f : M_1 \rightarrow M_2$ is K -quasiconformal for a constant $K > 1$ if and only if for any family Γ of C^1 Legendrian curves on M_1*

$$(3.9) \quad \frac{1}{K^2} \text{Mod}_{M_1}(\Gamma) \leq \text{Mod}_{M_2}(f(\Gamma)) \leq K^2 \text{Mod}_{M_1}(\Gamma),$$

where $f(\Gamma) = \{f(\gamma) \mid \gamma \in \Gamma\}$.

Proof. Assume $f : M_1 \rightarrow M_2$ is K -quasiconformal. Then f is contact, i.e., for contact forms η_1 and η_2 on M_1 and M_2 respectively, $f^*\eta_2 = \lambda \eta_1$ with a C^1 function λ on M_1 . Then

$$(3.10) \quad f^*(d\eta_2) = d\lambda \wedge \eta_1 + \lambda d\eta_1.$$

Therefore

$$(3.11) \quad f^*(d\eta_2|_{HM_2}) = \lambda d\eta_1|_{HM_1}.$$

For $j = 1, 2$, the Levi form L_j on M_j is a bilinear form on HM_j defined by

$$(3.12) \quad L_j(X, Y) = d\eta_j(X, J_j Y), \quad \text{for } X, Y \in HM_j.$$

By replacing η_j by $-\eta_j$, if necessary, we can always assume that L_j is positive definite. Hence L_j is a Hermitian form on HM_j . With respect to the Levi forms on M_1 and M_2 , we define

$$(3.13) \quad \begin{aligned} \lambda_1(q) &= \max_{Y \in H_q M_1, |Y|_1=1} |f_*(Y)|_2, \\ \lambda_2(q) &= \min_{Y \in H_q M_1, |Y|_1=1} |f_*(Y)|_2. \end{aligned}$$

Then (3.11) implies that $|\lambda| = \lambda_1 \lambda_2$. Moreover, (3.10) implies that

$$(3.14) \quad f^*(d\eta_2 \wedge \eta_2) = \lambda^2 d\eta_1 \wedge \eta_1.$$

Thus the Jacobian of f with respect to the volume forms $dv_1 = d\eta_1 \wedge \eta_1$ on M_1 and $dv_2 = d\eta_2 \wedge \eta_2$ on M_2 is $J(f) = (\lambda_1 \lambda_2)^2$.

For $\sigma_2 \in A(f(\Gamma))$,

$$(3.15) \quad \begin{aligned} \int_{f(\gamma)} \sigma_2 &= \int_{I_\gamma} |f_*(\gamma'(t))|_2 \sigma_2(f(\gamma(t))) dt \\ &\leq \int_{I_\gamma} \lambda_1 |\gamma'(t)|_1 \sigma_2(f(\gamma(t))) dt \\ &= \int_\gamma \lambda_1 \cdot \sigma_2 \circ f. \end{aligned}$$

Hence $\sigma_2 \in A(f(\Gamma))$ implies that $\lambda_1 \cdot \sigma_2 \circ f \in A(\Gamma)$.

On the other hand,

$$(3.16) \quad \begin{aligned} \int_{f(\gamma)} \sigma_2 &\geq \int_{I_\gamma} \lambda_2 |\gamma'(t)|_1 \sigma_2(f(\gamma(t))) dt \\ &= \int_\gamma \lambda_2 \cdot \sigma_2 \circ f. \end{aligned}$$

Hence $\sigma_1 \in A(\Gamma)$ implies $(\sigma_1/\lambda_2) \circ f^{-1} \in A(f(\Gamma))$. Therefore,

$$(3.17) \quad \begin{aligned} \text{Mod}_{M_2}(f(\Gamma)) &= \inf_{\sigma_2 \in A(f(\Gamma))} \int_{M_2} \sigma_2^4 dv_2 \\ &\leq \inf_{\sigma_1 \in A(\Gamma)} \int_{M_2} \left(\left(\frac{\sigma_1}{\lambda_2} \right) \circ f^{-1} \right)^4 dv_2 \\ &= \inf_{\sigma_1 \in A(\Gamma)} \int_{M_1} \frac{\sigma_1^4}{\lambda_2^4} \cdot (\lambda_1 \lambda_2)^2 dv_1 \\ &\leq K^2 \text{Mod}_{M_1}(\Gamma). \end{aligned}$$

$$\begin{aligned}
\text{Mod}_{M_1}(\Gamma) &= \inf_{\sigma_1 \in A(\Gamma)} \int_{M_1} \sigma_1^4 dv_1 \\
&\leq \inf_{\sigma_2 \in A(f(\Gamma))} \int_{M_1} (\lambda_1 \cdot \sigma_2 \circ f)^4 dv_1 \\
(3.18) \quad &= \inf_{\sigma_2 \in A(f(\Gamma))} \int_{M_2} \frac{(\lambda_1 \circ f^{-1} \cdot \sigma_2)^4}{(\lambda_1 \lambda_2)^2 \circ f^{-1}} dv_2 \\
&\leq K^2 \text{Mod}_{M_2}(f(\Gamma)).
\end{aligned}$$

Assume that f satisfies the inequalities (3.9). Then f must preserve the contact structures. Otherwise, there will be a point $q \in M_1$ with a tangent vector $X_q \in H_q M_1$ so that $f_*(X_q) \notin H_{f(q)} M_2$. f is C^1 , so there is a neighborhood U of q with a nonzero smooth section X of HM_1 on U so that $f_*(X_{q'})$ is not contact for each $q' \in U$. Let Γ be the family of trajectories of X , and shrink U appropriately such that $\text{Mod}_{M_1}(\Gamma) \neq 0$. But each curve in $f(\Gamma)$ is not Legendrian everywhere, hence not locally rectifiable. So $\text{Mod}_{M_2}(f(\Gamma)) = 0$ by Definition 3.1. Then f cannot satisfy (3.9) for this family Γ . This contradiction shows that f must be contact.

If f is not K -quasiconformal, there exists an open set $U \subset M_1$ so that $\lambda_1/\lambda_2 \geq K + \epsilon$, for some $\epsilon > 0$. Let X be the vector field on U so that $|X|_1 = 1$, $|f_*(X)|_2 = \lambda_2$. Let Γ be the family of trajectories of X in U . Then for any integrable function σ_2 on $f(U)$,

$$(3.19) \quad \int_{f(\Gamma)} \sigma_2 = \int_{\Gamma} \sigma_2 \circ f \cdot |f_*(X)|_2 = \int_{\Gamma} \lambda_2 \cdot \sigma_2 \circ f.$$

So $\sigma_2 \in A(f(\Gamma))$ if and only if $\sigma_1 \triangleq \lambda_2 \cdot \sigma_2 \circ f \in A(\Gamma)$. Hence

$$\begin{aligned}
\text{Mod}_{M_2}(f(\Gamma)) &= \inf_{\sigma_2 \in A(f(\Gamma))} \int_{M_2} \sigma_2^4 dv_2 \\
&= \inf_{\sigma_1 \in A(\Gamma)} \int_{f(U)} \left(\left(\frac{\sigma_1}{\lambda_2} \right) \circ f^{-1} \right)^4 dv_2 \\
(3.20) \quad &= \inf_{\sigma_1 \in A(\Gamma)} \int_U \left(\frac{\sigma_1}{\lambda_2} \right)^4 \cdot (\lambda_1 \lambda_2)^2 dv_1 \\
&= \inf_{\sigma_1 \in A(\Gamma)} \int_U \left(\frac{\lambda_1}{\lambda_2} \right)^2 \sigma_1^4 dv_1 \\
&\geq (K + \epsilon)^2 \text{Mod}_{M_1}(\Gamma).
\end{aligned}$$

Therefore, (3.9) implies that

$$(3.21) \quad (K + \epsilon)^2 \text{Mod}_{M_1}(\Gamma) \leq K^2 \text{Mod}_{M_1}(\Gamma).$$

But we can always choose U such that $\text{Mod}_{M_1}(\Gamma) \neq 0$. So (3.9) cannot be true for such Γ . Hence $\lambda_1/\lambda_2 \leq K$ on M_1 , i.e., f is K -quasiconformal. $\square$

We like to know if we can use the conclusion of Theorem 3.3 to define the quasiconformality for a homeomorphism $f : M_1 \rightarrow M_2$. The rest of this section gives some partial solutions to this problem.

Theorem 3.4. *If $f : M_1 \rightarrow M_2$ is a homeomorphism so that for a constant $K \geq 1$ and any curve family Γ which forms a smooth contact fibration of an open set in M_1*

$$(3.22) \quad \text{Mod}_{M_1}(\Gamma) \leq K^2 \text{Mod}_{M_2}(f(\Gamma)),$$

then f is ACL.

Proof. Let $U \subset M_1$ be an open set with a smooth contact fibration Γ , $X \neq 0$ be a horizontal vector field on U tangent to Γ . By replacing X by $X/|X|$, we can assume $|X| = 1$. Shrink U , if necessary, so that there is a smooth surface $S \subset U$ which intersects each fiber of Γ transversally once and only once. Parametrize fibers $\gamma \in \Gamma$ by t so that $X = \frac{\partial}{\partial t}$ and $\gamma(0) \in S$. $p : U \rightarrow S$ is the natural projection given by $\gamma(t) \mapsto \gamma(0)$.

Recall that Hermitian metrics on HM_1 and HM_2 can be extended canonically to Riemannian metrics on M_1 and M_2 respectively. Restricting the Riemannian metric on M_1 to S makes S a Riemannian 2-manifold. Let ω be the area form on S , and $A_\omega(E) \triangleq \int_E \omega$ the ω -area of a measurable set $E \subset S$. Define a set function F for measurable set $E \subset S$ by letting $F(E) = \text{vol}(f(p^{-1}(E)))$, where vol refers to the Riemannian volume on M_2 . Then Lebesgue's theorem (Theorem 23.5, [14]) asserts that F has finite derivatives at all points in S_1 with respect to ω -area, for a subset $S_1 \subset S$ with $A_\omega(S \setminus S_1) = 0$. We want to prove f is absolutely continuous along the fibers of Γ passing through points in S_1 .

For $q \in S_1$, let $D_r \subset S \cap U$ be a disc centered at q with radius r , $\gamma_q : I \rightarrow U$ be the fiber of γ passing through q . Take any sequence $t'_1, t''_1, t'_2, t''_2, \dots, t'_k, t''_k \in I$ such that

$$(3.23) \quad t'_1 < t''_1 < t'_2 < t''_2 \dots < t'_k < t''_k.$$

Let $\Delta t_j \triangleq t''_j - t'_j$. When $\max_{1 \leq j \leq k} \Delta t_j$ is small enough,

$$(3.24) \quad d_{r,j} \leq \Delta t_j \leq 2 d_{r,j},$$

where $d_{r,j}$ is the sub-Riemannian distance between the set $B'_{r,j} \triangleq \{\gamma(t'_j) \mid \gamma \in \Gamma, \gamma(0) \in D_r\}$ and the set $B''_{r,j} \triangleq \{\gamma(t''_j) \mid \gamma \in \Gamma, \gamma(0) \in D_r\}$. This is true because fibers of Γ are Legendrian and the geodesic with respect to the sub-Riemannian metric tangent to γ at $\gamma(t'_j)$ is locally a length minimizing curve (Theorem 5.4, [11]). For $j = 1, 2, \dots, k$, denote

$$R_{r,j} \triangleq \{\text{points } \gamma(t) \mid \gamma \in \Gamma, \gamma(0) \in D_r, t'_j \leq t \leq t''_j\}$$

and let

$$\Gamma_{r,j} \triangleq \{\text{curves } \gamma_j : t \mapsto \gamma(t) \mid \gamma \in \Gamma, \gamma(0) \in D_r, t'_j \leq t \leq t''_j\},$$

a contact fibration of $R_{r,j}$. Then by (3.22),

$$(3.25) \quad \text{Mod}_{M_1}(\Gamma_{r,j}) \leq K^2 \text{Mod}_{M_2}(f(\Gamma_{r,j})).$$

Next we use length-volume argument to give an estimate for $\text{Mod}_{M_1}(\Gamma_{r,j})$. For $\sigma \in A(\Gamma_{r,j})$ and $\gamma_j \in \Gamma_{r,j}$,

$$(3.26) \quad 1 \leq \left(\int_{\gamma_j} \sigma \right)^4 \leq \left(\int_{\gamma_j} \sigma^4 \right) \left(\int_{\gamma_j} 1 \right)^3 = (\Delta t_j)^3 \int_{\gamma_j} \sigma^4.$$

Integrating each term against ω over D_r , then

$$(3.27) \quad A_\omega(D_r) \leq c (\Delta t_j)^3 \int_{R_{r,j}} \sigma^4 dv_1.$$

where c is constant, dv_1 is the volume form of the Riemannian metric on M_1 . By (3.3) and (3.24),

$$(3.28) \quad A_\omega(D_r) \leq 8c d_{r,j}^3 \text{Mod}_{M_1}(\Gamma_{r,j}).$$

On the other hand, let $\delta_{r,j}$ be the sub-Riemannian distance between $f(B'_{r,j})$ and $f(B''_{r,j})$. Then $1/\delta_{r,j} \in A(f(\Gamma_{r,j}))$. Thus

$$(3.29) \quad \text{Mod}_{M_2}(f(\Gamma_{r,j})) \leq \frac{1}{\delta_{r,j}^4} \text{vol}(f(R_{r,j})).$$

Combining (3.25), (3.28) and (3.29), we have

$$(3.30) \quad (A_\omega(D_r))^{\frac{1}{4}} \delta_{r,j} \leq (8cK^2)^{\frac{1}{4}} d_{r,j}^{\frac{3}{4}} (\text{vol}(f(R_{r,j})))^{\frac{1}{4}}.$$

Summing (3.30) over j ,

$$(3.31) \quad \begin{aligned} & (A_\omega(D_r))^{\frac{1}{4}} \sum_{j=1}^k \delta_{r,j} \\ & \leq (8cK^2)^{\frac{1}{4}} \sum_{j=1}^k d_{r,j}^{\frac{3}{4}} (\text{vol}(f(R_{r,j})))^{\frac{1}{4}} \\ & \leq (8cK^2)^{\frac{1}{4}} \left(\sum_{j=1}^k d_{r,j} \right)^{\frac{3}{4}} \left(\sum_{j=1}^k \text{vol}(f(R_{r,j})) \right)^{\frac{1}{4}} \\ & \leq (8cK^2)^{\frac{1}{4}} \left(\sum_{j=1}^k d_{r,j} \right)^{\frac{3}{4}} (\text{vol}(f(p^{-1}(D_r))))^{\frac{1}{4}}. \end{aligned}$$

So by (3.24),

$$(3.32) \quad \sum_{j=1}^k \delta_{r,j} \leq (8cK^2)^{\frac{1}{4}} \left(\sum_{j=1}^k \Delta t_j \right)^{\frac{3}{4}} \left(\frac{F(D_r)}{A_\omega(D_r)} \right)^{\frac{1}{4}}.$$

Letting $r \rightarrow 0$,

$$(3.33) \quad \sum_{j=1}^k \delta_j \leq (8cK^2 F'(q))^{\frac{1}{4}} \left(\sum_{j=1}^k \Delta t_j \right)^{\frac{3}{4}},$$

where $\delta_j \triangleq \lim_{r \rightarrow 0} \delta_{r,j}$, i.e., the sub-Riemannian distance between $f(\gamma_q(t_j))$ and $f(\gamma_q(t_{j+1}))$. Hence f is absolutely continuous along γ_q , for $q \in S_1$. So f is ACL. $\square$

Corollary 3.5. *If a homeomorphism $f : M_1 \rightarrow M_2$ satisfies (3.22), then f has horizontal derivatives almost everywhere on M_1 in the sense that for any open set $U \subset M_1$ with a smooth section X of HM_1 on U and any smooth function on $f(U)$, $X(h \circ f)$ exists almost everywhere on U .*

4. EXTREMAL QUASICONFORMAL HOMEOMORPHISMS

In this section we will construct CR 3-manifolds M_1, M_2 and a quasiconformal diffeomorphism $f_0 : M_1 \rightarrow M_2$ such that $K(f_0) \leq K(f)$, for any C^2 quasiconformal homeomorphism $f : M_1 \rightarrow M_2$ which is homotopic to f_0 . M_1 to be constructed is the quotient of the 3-dimensional Heisenberg group over a lattice. So we start with the Heisenberg group.

The 3-dimensional Heisenberg group $\mathbb{H}^3$ is the space $\mathbb{R}^3$ endowed with the group structure defined by

$$(4.1) \quad (x, y, t)(u, v, s) = (x + u, y + v, t + s + 2yu - 2xv).$$

The standard contact structure on $\mathbb{H}^3$ is given by the contact form

$$(4.2) \quad \tilde{\eta} = -\frac{1}{2}ydx + \frac{1}{2}xdy + \frac{1}{4}dt.$$

The contact bundle $H\mathbb{H}^3 = \text{Ker } \tilde{\eta}$ has two global sections

$$(4.3) \quad \tilde{X} = \frac{\partial}{\partial x} + 2y\frac{\partial}{\partial t}, \quad \tilde{Y} = \frac{\partial}{\partial y} - 2x\frac{\partial}{\partial t}$$

which span $H\mathbb{H}^3$ everywhere and are invariant under the left group translation. The standard CR structure on $\mathbb{H}^3$ is

$$(4.4) \quad \tilde{J} : H\mathbb{H}^3 \rightarrow H\mathbb{H}^3, \quad \tilde{X} \mapsto \tilde{Y}, \quad \tilde{Y} \mapsto -\tilde{X}.$$

Note that $d\tilde{\eta} = dx \wedge dy$, thus the sub-Riemannian metric on $H\mathbb{H}^3$ determined by the area form $d\tilde{\eta}$ and the complex structure $\tilde{J}$, i.e., Levi form, is the one making $\{\tilde{X}, \tilde{Y}\}$ orthonormal. The canonical Riemannian extension of this metric is the Euclidean metric on $\mathbb{R}^3$.

Next we study the geodesics on $\mathbb{H}^3$ with respect to this sub-Riemannian metric. Since the metric is invariant under the left group translation, thus it suffices to study geodesics joining the origin $O = (0, 0, 0)$ and a generic point q . Denote by p the projection from $\mathbb{H}^3$ to the horizontal plane $P \triangleq \{t = 0\}$. It is easy to see that if γ is a rectifiable curve in $\mathbb{H}^3$ with respect to the sub-Riemannian metric, $p(\gamma)$ is rectifiable with respect to the Euclidean metric on $\{t = 0\}$, and the respective lengths of γ and $p(\gamma)$ coincide. The characterization of the geodesics given in the next theorem was first given by Korányi by studying Euler-Lagrange equations [3]. The following proof, which is due to Lempert [8], is a geometric one.

Proposition 4.1. *On $\mathbb{H}^3$, a minimal geodesic to connect the origin O and $q \in \mathbb{H}^3$ is on either a straight line if $q \in P$, or a helix whose projection to P is a circle if $q \notin P$.*

Proof. The conclusion is obvious if $q \in P$. Next we assume $q \notin P$. We first consider the case $p(q) = O$. i.e., $q = (0, 0, t)$ for some $t \neq 0$. Let

$$(4.5) \quad \gamma : [0, l] \rightarrow \mathbb{H}^3, \quad s \mapsto (x(s), y(s), t(s))$$

be any oriented rectifiable curve joining O and q . Then

$$(4.6) \quad -2yx' + 2xy' + t' = 0,$$

almost everywhere on γ by (4.2). Thus

$$(4.7) \quad t = \int_0^l t' ds = \int_{p(\gamma)} 2ydx - 2xdy = -4 \int_{\Omega} dx \wedge dy,$$

where Ω is the 2-chain on P so that $\partial\Omega = p(\gamma)$. Note length of γ with respect to the sub-Riemannian metric is equal to the length of $p(\gamma)$ on P with respect to Euclidean metric. Obviously, the length of $p(\gamma)$ is the minimal only when Ω is a simply connected domain with fixed area $|t|/4$. Furthermore, $p(\gamma)$ must be a circle if γ is a length minimizing geodesic joining O and q , by the isoperimetric property on the Euclidean plane.

If $p(q) \neq O$, the above reasoning with some slight modifications can be applied. For instance, Ω is a 2-chain bounded by the union of $p(\gamma)$ and the straight line segment from $p(q)$ to O . $\square$

Denote $A = (1, 0, 0), B = (\sigma, \tau, 0), C = (0, 0, 1) \in \mathbb{H}^3$ with $\tau \neq 0$. Define $\Gamma = \{n_1A + n_2B + n_3C \mid n_1, n_2, n_3 \in \mathbb{Z}\}$. The Euclidean translations by elements of Γ define a Γ -action. The contact structure and CR structure on $\mathbb{H}^3$ are invariant under this Γ -action. We consider the quotient space $M_1 \triangleq \mathbb{H}^3/\Gamma$. M_1 has a contact structure with the contact form η satisfying $\pi^*\eta = \tilde{\eta}$ and a CR structure J_1 inherited from $\tilde{J}$. Hence M_1 becomes a CR manifold. Let $\pi : \mathbb{H}^3 \rightarrow M_1$ denote the natural projection and $X = \pi_*\tilde{X}, Y = \pi_*\tilde{Y}$. Then with respect to the sub-Riemannian metric uniquely determined by η and J_1 , $\{X, Y\}$ is orthonormal. Note also $\bar{Z}_1 = \frac{1}{2}(X + iY) \in T^{0,1}M_1$. There is a canonical $\mathbb{R}$ -action on $\mathbb{H}^3$ defined by

$$(4.8) \quad (x, y, t) \mapsto (x, y, t + t'), \quad \text{for } t' \in \mathbb{R}.$$

This $\mathbb{R}$ -action is free and preserves both contact and CR structures on $\mathbb{H}^3$. The plane P can be regarded as the quotient space of this $\mathbb{R}$ -action. The natural projection $p : \mathbb{H}^3 \rightarrow P$ is isometric between the sub-Riemannian metric on $\mathbb{H}^3$ and the Euclidean metric on P . The free $\mathbb{R}$ -action on $\mathbb{H}^3$ commutes with the Γ -action. Thus it induces a free S^1 -actions on M_1 so that M_1 becomes a circle bundle over the torus $S_1 \triangleq \pi(P)$. The torus S_1 has a flat Riemannian metric inherited from the Euclidean metric on P . Denote the natural projection of this circle bundle by $\pi_1 : M_1 \rightarrow S_1$. Note the S^1 -action preserves the contact

and CR structures on M_1 . We call such a bundle $M_1 \xrightarrow{\pi_1} S_1$ a CR circle bundle over a flat torus.

For any constant $K \geq 1$, define a new CR structure M_2 on the contact structure of M_1 by declaring that

$$(4.9) \quad \overline{Z}_2 \triangleq \frac{1}{2}(\sqrt{K}X + \frac{i}{\sqrt{K}}Y) = \frac{K+1}{4\sqrt{K}}(\overline{Z}_1 + \frac{K-1}{K+1}Z_1)$$

is an $(0,1)$ tangent vector. Obviously, the identity mapping $f_0 : M_1 \rightarrow M_2$ is quasiconformal and $K(f_0)(q) = K$ for all $q \in M_1$.

Theorem 4.2. *Let $f : M_1 \rightarrow M_2$ be a C^2 homeomorphism homotopic to $f_0 : M_1 \rightarrow M_2$. Then $K(f) \geq K$.*

Note the CR structure on M_2 is also invariant under the circle action. So if S_2 denotes the same smooth torus as S_1 , but endowed with the complex structure induced from CR structure on M_2 , then $M_2 \xrightarrow{\pi} S_2$ is also a CR circle bundle.

Lemma 4.3 (Strichartz). *For any q in a sub-Riemannian manifold M , there is an $\epsilon > 0$ so that if $q_1 \in M$ with $d(q_1, q) \leq \epsilon$, there exists a length minimizing curve joining q_1 and q .*

This is Lemma 3.2 in [11], a proof was given there. If α is a curve, denote by $[\alpha]$ the homotopy class with fixed end points of α . Let $l(\alpha)$ be the length of α with respect to the sub-Riemannian metric on M .

Lemma 4.4. *Let $\alpha : [0, 1] \rightarrow M$ be a curve on a compact sub-Riemannian manifold to connect two points q_1 and q_2 , $\inf l(\beta)$ be taken over all $\beta \in [\alpha]$. Then this infimum is attained at a rectifiable curve $\tilde{\alpha} \in [\alpha]$.*

Proof. Let ϵ_q be the maximal $\epsilon > 0$ for $q \in M$ determined by Lemma 4.3. Since M is compact, $\inf_{q \in M} \epsilon_q > 0$. Take any r so that $0 < 3r < \inf_{q \in M} \epsilon_q$. Cover M with finitely many closed r -balls $B(q_j, r)$, $j = 1, 2, \dots, k$, with respect to the sub-Riemannian metric. Note any two points in the same $B(q_j, r)$ can be joined by a length minimizing curve within $B(q_j, 3r)$.

Let $L = \inf l(\beta)$ over all $\beta \in [\alpha]$. Let $\beta_n \in [\alpha]$ be curves such that $\lim_{n \rightarrow \infty} l(\beta_n) = L$. Divide each β by ordered points $p_{n0} = q_1, p_{n1}, p_{n2}, \dots, p_{nN} = q_2 \in \beta_n$ into ordered subcurves $\tau_{n1}, \tau_{n2}, \dots, \tau_{nN}$ so that each $\tau_{nj} \subset B(q_{m_{nj}}, r)$ for some integer m_{nj} with $1 \leq m_{nj} \leq k$. Note this N is uniform for all n by appropriately choosing β_n .

Let σ_{nj} be a length minimizing curve in $B(q_{m_{nj}}, 3r)$ to join the end points of σ_{nj} and $\sigma_n = \sigma_{n1}\sigma_{n2}\dots\sigma_{nN}$. Then since $B(q_{m_{nj}}, 3r)$ is simply connected, $\sigma_{nj} \in [\tau_{nj}]$, and hence $\sigma_n \in [\beta_n] = [\alpha]$. Furthermore

$$(4.10) \quad L \leq l(\sigma_n) \leq \sum_{j=1}^N l(\sigma_{nj}) \leq \sum_{j=1}^N l(\tau_{nj}) = l(\beta_n).$$

Therefore $\lim_{n \rightarrow \infty} l(\sigma_n) = L$.

Since M is compact, there is a sequence $\{n_j\} \subset \mathbb{N}$ so that for each $j = 1, 2, \dots, N-1$, p_{n_j} is convergent to some point p_j , when $n_j \rightarrow \infty$. Note $p_{j-1}, p_j \in B(q_{m_j}, r)$ for some integer m_j with $1 \leq m_j \leq k$. Connect p_{j-1}, p_j by a length minimizing curve $\alpha_j \subset B(q_{m_j}, 3r)$ and let $\tilde{\alpha} = \alpha_1 \alpha_2 \dots \alpha_N$. Then $\tilde{\alpha} \in [\alpha]$ and

$$(4.11) \quad l(\tilde{\alpha}) = \lim_{n_i \rightarrow \infty} l(\sigma_{n_i}) = L. \quad \square$$

The next two lemmas are the key properties to establish the extremality of f_0 described by Theorem 4.2.

Lemma 4.5. *The flow of diffeomorphisms h_t generated by X on M_1 preserve the volume form $dv_1 = d\eta \wedge \eta$.*

Proof. This is obvious since the flow of diffeomorphisms $\tilde{h}_s$ on $\mathbb{H}^3$ generated by $\tilde{X}$ are given by

$$(4.12) \quad \tilde{h}_s : (x, y, t) \mapsto (x + s, y, t + 2sy), \quad s \in \mathbb{R}$$

which preserve the volume form $d\tilde{\eta} \wedge \tilde{\eta} = dx \wedge dy \wedge dt$. $\square$

We define a sub-Riemannian metric on M_2 so that $\{\sqrt{K}X, \frac{Y}{\sqrt{K}}\}$ is orthonormal. Note this sub-Riemannian metric induces the same CR structure on M_2 . Denote the corresponding curve length by l_2 .

Lemma 4.6. *Let $\gamma : \mathbb{R} \rightarrow M_2$ be an integral line of the vector field $\sqrt{K}X$. Then for $[a, b] \subset \mathbb{R}$ and with respect to the sub-Riemannian metric on M_2 , the curve $\gamma|_{[a, b]}$ is the unique length minimizing curve in its homotopy class of curves with end points $\gamma(a)$ and $\gamma(b)$.*

Proof. Lifting the sub-Riemannian metric on M_2 to its universal covering $\mathbb{H}^3$, we obtain a new sub-Riemannian metric on $\mathbb{H}^3$. Let us call this new sub-Riemannian manifold $\mathbb{H}_2^3$. Then on $\mathbb{H}_2^3$, there is a characterization of geodesics similar to Proposition 4.1. The trajectories of $\sqrt{K}\tilde{X}$ are exactly those straight geodesics. Indeed, the sub-Riemannian metric is invariant under the $\mathbb{R}$ -action (4.8), hence a rectifiable curve γ in $\mathbb{H}_2^3$ has the same length with $p(\gamma)$ on P_2 . Here P_2 is the plane P endowed with the flat Riemannian metric so that $\sqrt{K}\frac{\partial}{\partial x}, \frac{1}{\sqrt{K}}\frac{\partial}{\partial y}$ are orthonormal. Therefore the x -axis, which is the trajectory of $\sqrt{K}\tilde{X}$ passing through O , is a geodesic without conjugate points on it. Note $\sqrt{K}\tilde{X}$ and the sub-Riemannian metric are invariant under the group left translation. Hence this lemma is true. $\square$

Proof of Theorem 4.2. Let $f : M_1 \rightarrow M_2$ be an C^2 quasiconformal homeomorphism. For $q \in M_1$, let $\gamma_{q,a} : [-a, a] \rightarrow M_1$ be the integral line of X so that $\gamma_{q,a}(0) = q$. Let $\Gamma_a = \{\gamma_{q,a} \mid q \in M_1\}$. Then by Theorem 3.4,

$$(4.13) \quad \text{Mod}_{M_1}(\Gamma_a) \leq K(f)^2 \text{Mod}_{M_2}(f(\Gamma_a)).$$

First note the length of $\gamma_{q,a}$ is $2a$, so $1/2a \in A(\Gamma_a)$ and

$$(4.14) \quad \text{Mod}_{M_1}(\Gamma_a) \leq \int_{M_1} \left(\frac{1}{2a}\right)^4 dv_1 = \left(\frac{1}{2a}\right)^4 \text{vol}(M_1).$$

For $\sigma \in A(\Gamma_a)$,

$$(4.15) \quad 1 \leq \left(\int_{\gamma_{q,a}} \sigma\right)^4 \leq \left(\int_{\gamma_{q,a}} 1\right)^3 \left(\int_{\gamma_{q,a}} \sigma^4\right) = (2a)^3 \int_{\gamma_{q,a}} \sigma^4.$$

That implies

$$(4.16) \quad \frac{1}{(2a)^3} \leq \int_{\gamma_{q,a}} \sigma^4 = \int_{-a}^a (\sigma(\gamma_{q,a}(s)))^4 ds.$$

Taking the integrals of both sides of (4.16) against dv_1 with respect to q over M_1 ,

$$(4.17) \quad \begin{aligned} \frac{1}{(2a)^3} \text{vol}(M_1) &\leq \int_{M_1} \left(\int_{-a}^a (\sigma(\gamma_{q,a}(s)))^4 ds \right) dv_1 \\ &= \int_{-a}^a \left(\int_{M_1} (\sigma(\gamma_{q,a}(s)))^4 dv_1 \right) ds \\ &= 2a \int_{M_1} \sigma^4 dv_1. \end{aligned}$$

The last equality in (4.17) is due to Lemma 4.5, which implies that

$$(4.18) \quad \int_{M_1} (\sigma(\gamma_{q,a}(s)))^4 dv_1 = \int_{M_1} \sigma^4 h_s^*(dv_1) = \int_{M_1} \sigma^4 dv_1$$

is independent of s . Therefore

$$(4.19) \quad \left(\frac{1}{2a}\right)^4 \text{vol}(M_1) \leq \int_{M_1} \sigma^4 dv_1, \quad \forall \sigma \in A(\Gamma_a).$$

Combining (4.14) and (4.19), we get

$$(4.20) \quad \text{Mod}_{M_1}(\Gamma_a) = \left(\frac{1}{2a}\right)^4 \text{vol}(M_1).$$

Next we estimate $\text{Mod}_{M_2}(f(\Gamma_a))$ in (4.13) for a quasiconformal homeomorphism f homotopic to $f_0 : M_1 \rightarrow M_2$. Since f is homotopic to f_0 , there is a continuous map $H : [0, 1] \times M_1 \rightarrow M_2$ so that

$$(4.21) \quad H(0, q) = f_0(q), \quad H(1, q) = f(q), \quad \forall q \in M_1.$$

Let $\alpha_q : [0, 1] \rightarrow M_2$ be a curve given by $t \mapsto H(t, q)$. Define $G(q) = \inf l_2(\beta)$ over all $\beta \in [\alpha_q]$. Then by Lemma 4.4, $G(q)$ is attained at a rectifiable curve $\tilde{\alpha}_q \in [\alpha_q]$.

Now we prove that $G(q)$ is continuous on M_1 . When q_1 is close enough to q on M_1 , $f_0(q_1)$ is close enough to $f_0(q)$ so that there exist a length minimizing curve δ_1 joining $f_0(q)$, $f_0(q_1)$ according to Lemma 4.3. Let $\delta_2 = f(\delta_1)$. Then

$$(4.22) \quad \delta_1 \tilde{\alpha}_{q_1} \delta_2^{-1} \in [\alpha_q], \quad \delta_1^{-1} \tilde{\alpha}_q \delta_2 \in [\alpha_{q_1}].$$

This implies

$$(4.23) \quad \begin{aligned} l_2(\tilde{\alpha}_q) &\leq l_2(\delta_1) + l_2(\tilde{\alpha}_{q_1}) + l_2(\delta_2^{-1}), \\ l_2(\tilde{\alpha}_{q_1}) &\leq l_2(\delta_1^{-1}) + l_2(\tilde{\alpha}_q) + l_2(\delta_2). \end{aligned}$$

In other words,

$$(4.24) \quad |G(q_1) - G(q)| \leq l_2(\delta_1) + l_2(\delta_2).$$

Then there exists $A > 0$ so that

$$(4.25) \quad G(q) \leq A < \infty, \quad \forall q \in M_1,$$

since M_1 is compact.

For a curve $\gamma : [a, b] \rightarrow M_1$, $\tilde{\alpha}_{\gamma(a)} f(\gamma) \tilde{\alpha}_{\gamma(b)}^{-1} \in [f_0(\gamma)]$. Then by Lemma 4.6 and (4.25),

$$(4.26) \quad l_2(f_0(\gamma)) \leq l_2(\tilde{\alpha}_{\gamma(a)}) + l_2(f(\gamma)) + l_2(\tilde{\alpha}_{\gamma(b)}^{-1}) \leq 2A + l_2(f(\gamma)).$$

Applying (4.26) to $\gamma = \gamma_{q,a} \in \Gamma_a$, we obtain

$$(4.27) \quad 2\sqrt{K}a \leq 2A + l_2(f(\gamma_{q,a})).$$

Hence $1/(2\sqrt{K}a - 2A) \in A(f(\Gamma_a))$. So we have the following estimate.

$$(4.28) \quad \text{Mod}_{M_2}(f(\Gamma_a)) \leq \int_{M_2} \left(\frac{1}{2\sqrt{K}a - 2A} \right)^4 dv_2 = \left(\frac{1}{2\sqrt{K}a - 2A} \right)^4 \text{vol}(M_2).$$

Note $\text{vol}(M_1) = \text{vol}(M_2) \neq 0$, so by (4.13), (4.20) and (4.28), we have

$$(4.29) \quad \left(\sqrt{K} - \frac{A}{a} \right)^2 \leq K(f).$$

Letting $a \rightarrow \infty$, we get that $K \leq K(f)$. $\square$

Remark. In horizontal directions, the extremal quasiconformal homeomorphism f_0 behaves as a stretching by the constant factor $\sqrt{K}$ along trajectories of X and a compressing by the same factor along trajectories of JX . The generator T of the circle action is transversal to the contact bundle. In this transversal direction T , f_0 is equivariant under the circle action since it is simply the identity mapping while M_1 and M_2 have the same circle action.

REFERENCES

1. Bers, Lipman, *Quasiconformal mappings and Teichmüller's theorem*, Analytic functions, edited by R. Nevanlinna et al. (1960), Princeton University Press, Princeton, New Jewsey, 89–119.
2. Hamenstädt, Ursula, *Some regularity theorems for Carnot-Caratheodory metrics*, J. Differential Geometry **32** (1990), 819–850.
3. Korányi, A., *Geometric aspects of analysis on the Heisenberg group*, Topics in Modern Harmonic Analysis (1983), Istituto Naziiionale di Alta Matematica, Roma.
4. Korányi, A. and Reimann, H.M., *Foundations for the theory of quasiconformal mappings on the Heisenberg group*, Advances in Math (to appear).
5. ———, *Quasiconformal mappings on CR manifolds*, Conference in honor of E. Vesentini, Springer Lecture Notes, no. 1422 (1988), 59–75.
6. ———, *Quasiconformal mappings on the Heisenberg group*, Inventiones Mathematicae **80** (1985), 309–338.
7. Lempert, László, *On three dimensional Cauchy-Riemann Manifolds*, Journal of AMS **5 no.4** (1992), 923–969.
8. ———, *Private communication*.
9. Li, Zhong, *Quasiconformal mappings and their applications in the theory of Riemann surfaces*, Science Publisher, Beijing, 1988.
10. Pinchuk, S.I. and Tsyganov, Sh.I., *The smoothness of CR-mappings between strictly pseudoconvex hypersurfaces*, Math. USSR Izvestiya **35 no.2** (1990), AMS, 457–467.
11. Strichartz, Robert S., *Sub-Riemannian geometry*, J. of Differential Geometry **24** (1986), 221–263.
12. Tang, Puqi, *Quasiconformal homeomorphisms on CR manifolds with symmetries*, preprint (1993), 1–20.
13. Teichmüller, Oswald, *Extremale quasikonforme Abbildungen und quadratische Differentiale*, Abh. Prs. Akad. Wiss., math.-naturw. Kl. 22 (1939), 1–197, or Gesammelte Abhandlungen - Collected Papers, edited by L.V. Ahlfors and F.W. Gehring (1982), Springer-Verlag, New York Berlin, 335–531.
14. Väisälä, J., *Lectures on n-dimensional Quasiconformal Mappings*, Lecture Notes in Mathematics, no. 229, Springer-Verlag, New York Berlin, 1971.

DEPARTMENT OF MATHEMATICS, PURDUE UNIVERSITY, WEST LAFAYETTE, IN 47907, USA

E-mail address: tang@math.purdue.edu

Version 12.04.1993. Run at euclid.math.purdue.edu